CALMAGINE SOLUTION

Peripheral analgesic antipyretic and smooth-musculature antispasmodic

Composition

Solution for injection :
Dipyrone.....50 g
Benzyl alcohol..... 1 ml
Excipient to..... 100 ml

Properties

Derived from Aminophenazone (pyrazoled core), Noramidopyrine (or Dipyrone) is an excellent pheripheral analgesic. It has also antipyretic properties combined with antispasmodic action on smooth muscle fibres.

Indications

In cattle, pigs, horses, dogs and cats : antispasmodic.

Administration and dosage

Deep intramuscular and intravenous administrations.
Horses, pigs : 5 - 10 ml per 100 kg of b.w. (2.5 - 5 g of Dipyrone per 100 kg of b.w.).
Cattle : 5 ml per 100 kg of b.w.(2.5 g of Dipyrone per 100 kg of b.w.).
Dogs, cats : 1 ml per 2.5 - 5 kg of b.w. (100 - 200 mg of Dipyrone per kg of b.w.).
The effect of Calmagine® begins 5 - 10 minutes after the injection to persist for approximately 1 hour.
The injection may be repeated if the pain and agitation of the patient recur.

Special warnings

During intramuscular injection, it is recommended to distribute the total dose at several points.
Subcutaneous irritation may be produced by the injection of the solution, although necrosis and suppuration do not occur. The intramuscular injections should thus be carried out under strict aseptic conditions.

Contra-indications

Do not use in conjunction with barbiturates, Phenylbutazone or Chlorpromazine.
In the absence of withdrawal period for milk, do not use in dairy, lactating or in period of drying up females producing milk for consumption nor in pregnant females producing milk for future consumption, two months before parturition.

Withdrawal period

Meat and offal :
- Cattle : 18 days,
- Horses, pigs : 15 days.

Milk : refer to section "Contra-indications".

Category

List I.
Prescription only medicine. This drug should be stored during duration of its withdrawal period.

Presentation

100 ml bottle
M.A. N°688 579.0 from the 06/22/92